arXiv:1409.7903v1 [math.GR] 28 Sep 2014

The Answers to a Problem and Two Conjectures about OD-Characterization of Finite Groups

Ali Mahmoudifar & Behrooz Khosravi
Dept. of Pure Math., Faculty of Math. and Computer Sci.,
Amirkabir University of Technology (Tehran Polytechnic),
424, Hafez Ave., Tehran 15914, IRAN,
e-mail: khosravibbb@yahoo.com
alimahmoudifar@gmail.com

Abstract

In [Akbari and Moghaddamfar, Recognizing by order and degree pattern of some projective special linear groups, *Internat. J. Algebra Comput.*, 2012] the authors possed the following problem:

Problem. *Is there a simple group which is k-fold OD-characterizable for $k \geq 3$?*

In this paper as the main result we give positive answer to the above problem and we introduce two simple groups which are k-fold OD-characterizable such that $k \geq 6$.

Also in [R. Kogani-Moghadam and A. R. Moghaddamfar, Groups with the same order and degree pattern, *Science China Mathematics*, 2012], the authors possed two conjectures as follows:

Conjecture 1. *All alternating groups A_m with $m \neq 10$ are OD-characterizable.*

Conjecture 2. *All symmetric groups S_m, with $m \neq 10$, are n-fold OD-characterizable, where $n \in \{1, 3\}$.*

In this paper we find some alternating and some symmetric groups such that these conjectures are not true for them.

2000 AMS Subject Classification: 20D05, 20D60, 20D08.
Keywords : Alternating group, prime graph, degree pattern, order, characterization.

1 Introduction

In this paper every group is finite. If n is a natural number, then we denote by $\pi(n)$, the set of all prime divisors of n. If G is a finite group, then $\pi(|G|)$ is denoted by $\pi(G)$. The prime graph $GK(G)$ of a group G is defined as a graph with vertex set $\pi(G)$ in which two distinct primes $p, q \in \pi(G)$ are adjacent (and we write $p \sim q$) if G contains an element of order pq. In [5, Proposition 1.1], it is proved that if $r, s \in \pi(A_n) \setminus \{2\}$, then $r \nsim s$ in $GK(A_n)$ if and only if $r + s > n$ and $2 \nsim r$ in $GK(A_n)$ if and only if $r + 4 > n$.

Definition 1.1. ([3]) *Let G be a group and $\pi(G) = \{p_1, p_2, \ldots, p_k\}$ where $p_1 < p_2 < \cdots < p_k$. Then the degree pattern of G is defined as follows:*

$$D(G) := (\deg(p_1), \deg(p_2), \ldots, \deg(p_k)),$$

where $\deg(p_i)$, $1 \leq i \leq k$, is the degree of vertex p_i in $GK(G)$.

For a group G, denote by $h_{OD}(G)$ the number of nonisomorphism finite groups H, such that $|H| = |G|$ and $D(H) = D(G)$.

Definition 1.2. ([3]) *Let G be a group. We call that G is k-fold OD-characterizable if $h_{OD}(G) = k$. When $k = 1$, the finite group G is called OD-characterizable.*

OD-characterization of some finite groups are considered by some authors (see the references of [3]). In [4], the following problem about OD-characterization of finite simple groups is possed (see also [1, 3, 4]):

Problem 1.3. *Is there a simple group which is k-fold OD-characterizable for $k \geq 3$?*

Also in [3], the authors put the following conjectures:

Conjecture 1.4. *All alternating groups A_m with $m \neq 10$ are OD-characterizable.*

Conjecture 1.5. *All symmetric groups S_m, with $m \neq 10$, are n-fold OD-characterizable, where $n \in \{1, 3\}$.*

Until now, Conjecture 1.4 is valid for $n = p$, $p+1$ and $p+2$, where p is an odd prime number.

In this paper as the main result we prove the following theorem and using this theorem we introduce some finite simple groups which are k-fold OD-characterizable, where $k \geq 6$. Also we present some counterexamples for the above conjectures.

Main Theorem 1.6. *Let p be an odd prime number such that $p + 2$ and $p + 4$ are not prime and $p + 6 = 5^\alpha$, for some natural number α. Also let H and T be some groups such that $|H| = p + 6 = 5^\alpha$ and $|T| = 2(p + 6)$. Then we have:*

(1) $|A_{p+6}| = |A_{p+5} \times H|$ *and* $D(A_{p+6}) = D(A_{p+5} \times H)$,

(2) $|S_{p+6}| = |S_{p+5} \times H|$ *and* $D(S_{p+6}) = D(S_{p+5} \times H)$,

(3) $|S_{p+6}| = |A_{p+5} \times T|$ *and* $D(S_{p+6}) = D(A_{p+5} \times T)$.

Remark 1.7. *Easily we can see that $p = 619 = 5^4 - 6$, $15619 = 5^6 - 6$, $9765619 = 5^{10} - 6$ satisfy the hypothesis of the main theorem.*

Corollary 1.8. *If $(G, K) = (A_{625}, A_{624})$ or (S_{625}, S_{624}), then the following groups have the same order and degree pattern: G, $K \times Z_{625}$, $K \times Z_{125} \times Z_5$, $K \times Z_{25} \times Z_{25}$, $K \times Z_{25} \times Z_5 \times Z_5$, $K \times Z_5 \times Z_5 \times Z_5 \times Z_5$.*

Corollary 1.9. *We have $h_{OD}(A_{625}) \geq 6$. In Particular, the simple group A_{625} is an answer for Problem* 1.3 *and also we see that Conjecture* 1.4 *is not true.*

Corollary 1.10. *We have $h_{OD}(S_{625}) \geq 6$. In Particular, Conjecture* 1.5 *is not true.*

Corollary 1.11. *Let $p = 5^\alpha - 6$ satisfies the assumptions of the main theorem. If m is the number of nonisomorphic finite groups of order 5^α, then $h_{OD}(A_{p+6}) \geq m + 1$ and $h_{OD}(S_{p+6}) \geq m + 1$.*

2 Proof of The Main Theorem

Let p be a prime number such that $p+2$ and $p+4$ are not prime and $p+6=5^{\alpha}$, for some natural number α. The proof of the main theorem for symmetric groups is similar to alternating groups.

First we show that $GK(A_{p+6}) = GK(A_{p+5})$. By assumption, $p+6$ is a power of 5 and so $\pi(A_{p+6}) = \pi(A_{p+5})$. Since $p+2$ and $p+4$ are not prime numbers, by [5, Proposition 1.1], we get that every vertex is adjacent to 2, 3 and 5 in $GK(A_{p+5})$ and $GK(A_{p+6})$. Then it is sufficient to check the adjacency of odd primes in these graphs.

Let $r, s \in \pi(A_{p+5}) \setminus \{2\}$. By [5, Proposition 1.1], if $r \sim s$ in $GK(A_{p+5})$, then $r+s \leq p+5 < p+6$ and so $r \sim s$ in $GK(A_{p+6})$. Let $r \nsim s$, in $GK(A_{p+5})$. Then by [5, Proposition 1.1], we have $r+s > p+5$. Since $p+6$ is odd, we get that $r+s \neq p+6$, and so $r+s > p+6$. This implies that $r \nsim s$ in $GK(A_{p+6})$.

Let $r, s \in \pi(A_{p+6}) \setminus \{2\}$. If $r \nsim s$ in $GK(A_{p+6})$, then by [5, Proposition 1.1], $r+s > p+6 > p+5$ and so $r \nsim s$ in $GK(A_{p+5})$. Since $r+s \neq p+6$, if $r \sim s$ in $GK(A_{p+6})$, then $r+s \leq p+5$ and so $r \sim s$ in $GK(A_{p+5})$.

Therefore by the above discussion we conclude that $GK(A_{p+6}) = GK(A_{p+5})$. Now if H is a group such that $|H| = p+6 = 5^{\alpha}$, then $|A_{p+6}| = |A_{p+5} \times H|$. Also since 5 is adjacent to each vertex in $GK(A_{p+6})$ and $|H| = 5^{\alpha}$, then we get that $GK(A_{p+6}) = GK(A_{p+5} \times H)$. Therefore for every group H, such that $|H| = p+6 = 5^{\alpha}$, we have $|A_{p+6}| = |A_{p+5} \times H|$ and $D(A_{p+6}) = D(A_{p+5} \times H)$.

References

[1] B. Akbari, A. R. Moghaddamfar, Recognizing by order and degree pattern of some projective special linear groups, *Internat. J. Algebra Comput.* **22 (6)**, (2012) 1250051, 22 pp.

[2] A. A. Hoseini, A. R. Moghaddamfar, Recognizing alternating groups A_{p+3} for certain primes p by their orders and degree patterns, *Front Math. China,* **5 (3)**, (2010) 541–553.

[3] R. Kogani-Moghadam and A. R. Moghaddamfar, Groups with the same order and degree pattern, *Science China Mathematics,* **55 (4)**, (2012) 701–720.

[4] A. R. Moghaddamfar; A. R. Zokayi, Recognizing finite groups through order and degree pattern, *Algebra Colloq.*,**15 (3)**, (2008) 449-456.

[5] A. V. Vasil'ev and E. P. Vdovin, An adjacency criterion for the prime graph of a finite simple group, *Algebra Logic,* **44 (6)**, (2005) 381–406.